

Se-ACQUISITION AND REACTIVE OXYGEN SPECIES ROLE IN GROWTH, PHOTOSYNTHESIS, PHOTOSYNTHETIC PIGMENTS, AND BIOCHEMICAL CHANGES IN ESSENTIAL OIL(S) MONOTERPENE OF GERANIUM (*PELARGONIUM GRAVEOLENS* L.HER.' EX. AIT.)

A.Misra, A.K. Srivastava, N.K. Srivastava and A. Khan

Central Institute of Medicinal and Aromatic Plants, P.O.CIMAP, Kukrail Picnic Spot Road Lucknow – 226015, India.

Abstract: Se is an essential element required for various metabolic pathways and act as an antioxidant in the various redox-reactions of primary and secondary plant production of – biomolecules. Geranium is an important essential monoterpene oil(s) bearing plant. Culturing the plant at different doses of Se from 0-1.0 $\mu\text{g Se ml}^{-1}$ revealed that Se plays an important role as an antioxidant promoter, apart from its micronutrient essentiality. 0.25 $\mu\text{g Se ml}^{-1}$ is the critical concentrations for maximum content of (0.21%) total essential monoterpene oil(s). At concentration below and above 0.25 $\mu\text{g Se ml}^{-1}$, the CO_2 assimilation rate, photosynthetic pigments content and ultimately the accumulation of essential monoterpene oil(s) are affected. The maximum peroxidase and SOD activities were obtained at 0.25 $\mu\text{g Se ml}^{-1}$, with the production of biomolecule geraniol. Results revealed an oxidoreducible reaction of Se in the formation of monoterpene essential oil(s) and possibly for the major constituents Geraniol.

Key words: Se-acquisition, reactive oxygen species, photosynthesis, essential oil(s) monoterpene, *Pelargonium graveolens*.

INTRODUCTION

Selenium is a trace element that can function as an essential nutrient for humans and animals or as an environmental toxicant; the boundary between the two is narrow and depends on its chemical form, concentration, and other environmentally regulating variables (Fan et al., 2002; Shardendu et al., 2003). Further, it is an important microelement, exists in small amounts in microorganisms, plants, animals and humans. Although selenium is an essential trace nutrient important to humans and most other animals as an antioxidant, toxicity occurs at high concentrations due to replacement of sulphur with selenium in amino acids resulting in incorrect folding of the protein and consequently nonfunctional proteins and enzymes. Distribution of selenium and its species in plants .It was shown, that pea was good accumulators of selenium. The selenium content of pea seeds obtained from the untreated (UT group), once (OT) and twice (TT) foliarly treated plants was determined. The selenium content of pea seeds obtained from the untreated, once and twice foliarly treated plants was, in each case, directly proportional to the number of spraying applications. Seeds are usually a moderate source of selenium, but several studies dealing with cereal and legume seeds showed, that they are able to accumulate high amounts of selenium (Stadlober et al., 2001; Smrkolj et al., 2005; Smrkolj et al., 2006a). Higher selenium contents were found in leaves than in stems of plants grown from both OT and TT seeds, and these contents were, as for the seeds, directly proportional to the number of original foliar treatments (Smrkolj et al., 2006a). Selenomethionine was the only selenium compound found in supernatants by anion and cation exchange chromatography, comprising 49 and 67 % of the total selenium content in the OT and TT groups, respectively (Smrkolj et al., 2006a). The results of the study show that selenium enriched pea seeds are potential source of dietary selenium, on account of their ability to accumulate Se, and that this selenium is present mainly as SeMet, known to be favourable for human consumption (Smrkolj et al., 2006a). Common (*Fagopyrum esculentum*) and tartary (*Fagopyrum tataricum*) buckwheat was treated by spraying the leaves with a water solution containing in the form of sodium selenate in the flowering period. The selenium content in all parts of plant was found to be less than 200 ng g⁻¹ in non-treated and in the range 2700–4650 ng g⁻¹ in selenium treated buckwheat. Exposure to UV-B radiation lead to higher selenium accumulation in flowers of both selenium enriched cultivars. In flowers and leaves, on average 11% of the selenium content was soluble and in the form of Se(VI), representing between 0.6% (flowers) and 3% (leaves) of the selenium content.

Light stress can promote an accumulation of active oxygen species(AOS) in the chloroplasts in situation where the antioxidant capacity to detoxify AOS is exceeded. At low temperature the risk of over –excitation of photosynthesis and formation of AOS is increased as many energy –requiring processes are slowed down and the balance between energy supply and demand is lost resulting in reduction of oxygen (4-6) . Antioxidant capacity increases during cold acclimation in several plants as an adaptive mechanism to low temperature . enhanced antioxidant capacity is also evident during acclimation to photoinhibition indicating the important role of AOS scavenging capacity for preventing the injury to the photosynthetic apparatus(9). Selenium is an essential micronutrient needed in antioxidation and hormone balance in plants(10-12). Recent findings show that Se is toxic at higher concentrations , it can exert beneficial effects on plants at low concentrations. Se can increase the tolerance of plants to UV – induced oxidative stress as well as delay in senescence and promote the growth of aging(13-15). The antiaging effect was related to decreased lipid peroxidation and enhanced glutathione peroxidase activity in Se-treated plants. In plants grown at higher light intensities Se counteracted senescence-related oxidative stress and maintain the green leaf colour longer. Se also decrease the activity of Superoxide dismutase and lowered the amount of tocopherols . Se also promotes H₂O₂ scavenging by increasing the glutathione peroxidase activity and by the side enhances spontaneous disproportion of super oxide radicals to H₂O₂ and thus decreases the need for high superoxide dismutase activity(15). One approach to improve the crop plant tolerance to environmental stresses is to increase their antioxidant capacity(16,17). The most recent studies on Zea mays have shown that elevated antioxidant levels can protect the photosynthetic apparatus from oxidative damage. (18,19). In Geranium which is to be cold loving plant improved photosynthetic capacity and productivity at low temperature could be achieved by enhancing the detoxification of AOS. Study was undertaken to determine whether Se contributes to recovery of photosynthesis from prolonged high light stress in Geranium. The hypothesis was that through its antioxidative effect Se can alleviate oxidative stress in chloroplast. Further, the response of Geranium to Se supplementation were investigated by monitoring chlorophyll formation its fluorescence and the transcription of antioxidative enzymes. In heavy manuring - N,P,K and excessive vermiculturing as an organic fertigation to increase the crop productivity and sustainability leads, to the major environmental hazardous and limiting factors in crop production .

The general response of plants to increase the ion concentration, includes osmotic stress, specific ion toxicity and further the nutrient deficiencies, thus affecting a range of physiological processes involved in cell metabolism (Munns, 2002). Further at higher altitudes cultivation, the low pH and high Fe toxicity, apart from heavy metal toxicity due to vermiculturing in plains affects the productivity and biomass of the plants. These cumulative effects of salts complexes and heavy metal in roots, including the other environmental stresses, are known to be mediated at least partially, by an advocative enhanced generation of Reactive Oxygen Species (ROS) and free radicals. ROS such as hydrogen peroxide (H_2O_2), are generated from the autoxidation of lipids and H_2O_2 production through water-water cycle (Asada 1999, Havaux et al. 2003). Leaf antioxidant systems include the enzymes and antioxidant metabolites of the so called water-water cycle, which functions to scavenge superoxide that is produced from the reduction of oxygen by PSI (Asada, 1999). In the water-water cycle, superoxide dismutase (SOD) catalyses the conversion of superoxide to hydrogen peroxide, which is then converted to water via the enzyme ascorbate peroxidase using the ascorbate as a reductant. Formations of these ROS by strong sun shine and UV irradiations are likely to damage several cellular components such as lipids, proteins, nucleic acid and DNAs through oxidation (Singh 1989, Sawa 2003). Plants contain a wide variety of chemicals that have potent antioxidants – B-carotene, L-tocopherol, ascorbates, Vit.E, Cr, Zn, and Se metals which effects overall metabolism and physiology of the plants through antioxidants activities in scavenging the free-radicals formed in stresses caused by intrinsic, extrinsic biotic and abiotic factors. Stresses causes a severe imbalance between light absorption and its utilization via photosynthesis. In such conditions, plants employ photoreceptive mechanisms to deal with excess light absorbed by chlorophyll. Two key mechanisms operates: one Xanthophyll cycle-dependent thermal energy dissipation of excess light within the light-harvesting complexes; and second leaf antioxidants enzymes and metabolites which scavenges free radicals and dangerous ROS that are produced in excess stresses and in the presence of excess light. (Verhoeven et al. 2005). Stresses of light harvesting system of higher light illumination light intensity involves major increased in xanthophyll – cycle dependent energy dissipation utilizes xanthophylls carotenoids in a presence of high light energy absorbed by chlorophyll is converted to heat and dissipated harmlessly, thus preventing the formation of singlet oxygen within the light harvesting complexes (Deoimmming – Adams and Adms, 1996). Thus, increases include increased pool of xanthophylls – cycle pigments during stresses of light and thus increased in the allocation of absorbed light towards energy dissipation.

Under stresses leaf antioxidants system include the enzyme and antioxidants and its metabolites of the so called water-water cycle, which scavange superoxides that is produced from the reduction of oxygen by PS1 (Asada, 1999). Here, the superoxide dismutase(SOD) catalyzes the conversion of superoxide to hydrogen peroxide(H_2O_2), which then converted to water via ascorbate peroxidases using the ascorbate as a reductants. Excess H_2O_2 further affects carbon reduction through Calvin cycle(Takeda et al.1995) in chloroplast. L-tocopherol is a lipid – soluble antioxidant metabolite that exists in thalokoid membrane and functions in the detoxification of lipid radicals formed in the presence of singlet oxygen(Havaux, 2003).

Stimulating effects to the additions of small amount of Se on plant growth and yield have been observed by several researchers (Adel M Zayed and Terry, 2003; Terry, 1981). No conclusive evidence yet of an essential role of Se in plant metabolisms but to effect the growth and productivity of plants at low concentration utilization (Huffman and Allaway,1973), either by solubilization of Fe^{+3} to Fe^{+2} active Fe for plant uptake or in antioxidant activities for defence mechanisms in controlling ROS. Therefore the acquisition of Se on growth and physiology of Geranium with antioxidant activities will be studied in detail.

Geranium, *Pelargonium graveolens* L. `Her, ex Ait. Syn., *P. roseum* Willd of family Geraniaceae is the only source of one of the most important essential monoterpene oil(s) called the oil of geranium. It is commonly known as rose geranium (Anonymous, 1966, 1975, 1982 and Erickson, 1976). It is distinctly different from the horticultural Geranium, which are basically the ornamental and have no commercial usage in perfumery industries (Douglas, 1969). *P. graveolens* widely grown varieties are Algerian/Funion, Kelkar/Egyption and Bourbon/Reunion (Rajeshwar Rao and Bhattacharya 1992) is popularly cultivated in India. Steam distillation of the shoot biomass of Geranium yields the geraniol and citronellol rich monoterpene oil(s), extensively used for perfuming soaps, and cosmetics to which it imparts a pronounced and lasting rose odour, for high grade perfumes. It is largely used in flavoring tobacco products, tooth pastes, ointments and other pharmaceutical preparations (Ranade, 1988).

Essential oil biosynthesis in Geranium is strongly influenced by Se-acquisition and the stresses caused by Se on nutrition and growth. Se is involved in carbon assimilation and saccharide accumulation, free radical removal, in antioxidant enzymes and role in carbon utilized in terpene biosynthesis and the overall growth of the plants. Furthermore, the role of Se in influencing essential monoterpene oil(s) accumulation seems particularly important.

The requirement of Zn antioxidant for Japanese mint and Pelargonium and its limitations imposed on photosynthetic carbon metabolism and translocation in relation to essential oil accumulation in mint have been documented by Misra and Sharma (1991) and Misra et al. (2004), respectively. While Se as an antioxidants enzymes for free radical quenching in Geranium have not fully been documented and not much work has been done so far.

In the present paper we report on the role of – Se, as a stimulant of free-radicals quenching through Se affected antioxidants enzyme activity, in Geranium subjected to Se stress. Simultaneously, photosynthetic efficiency in terms of changes in CO₂ exchange rate, photosynthetic pigments, leaf fresh mass, dry mass, leaf area, Se content in whole plant shoot biomass and oil yield were also determined.

Materials and Methods:

Plant: Plant tips (5-6 inches) with 3-4 leaves of Geranium (*Pelargonium graveolens*) of Bourbon genotype were obtained from the farm nursery of the CIMAP, Lucknow, India. Uniform cuttings were initially planted in 10000 cm³ earthen pots filled with purified silica sand (Agarwala and Sharma, 1961), for the development of roots. After 15 days, rooted cuttings were transferred to 2500 cm³ pots. The salts used in nutrient solution culture were purified for Zn (Hewitt, 1952). Hoagland and Arnon's (1959) nutrient solution was used in the experiment except Fe which was supplied as Fe-EDTA. Three pots each of Zn treatments ranging from 0.0 to 1.0 µg Zn ml⁻¹ were maintained in controlled glass house condition at ambient temperature (30±5°C) and irradiance (between 800-1000 µmol m⁻² s⁻¹). The nutrient solution in each treatment was added at alternate days. With onset of deficiency and toxicity (after 20 days) growth observation and detailed physiological and biochemical data with growth attributes were performed.

CO₂ exchange rate (P_N): CO₂ exchange rate was measured using a computerized portable photosynthesis system (Srivastava and Misra, 1991) (Model LiCOR 6000, LiCOR, USA).

Chlorophyll (Chl) content: A known mass of leaf tissue (3rd leaf) was extracted with 80% acetone and absorbance was recorded on spectrophotometer (Pye Unicam PU8610, USA) for determination of Chl a and b according to Arnon (1949)., and Carotenoids were calculated as described by Deming-Adams(1992).

Growth attributes and Zn analysis: Leaf fresh and shoot dry mass and area (area meter LICOR Li-3000) were recorded. For tissue elemental analysis 1.g. dried leaf samples were digested with IN HCl at 60°C for 24 h. Aliquot samples of the clear digest were diluted with water (10 cm³) and analyzed for Zn by atomic absorption spectrophotometer (Pye Unicam SP 2800) (Misra and Sharma 1991). Antioxidant reactive peroxidase enzyme activity was estimated as described earlier (Shanon *et al.*, 1960). Using 2 g of fresh chopped leaves at position 3rd, were homogenized with 5 ml of 0.1 M phosphate buffer (pH 6.8). Each treatment was replicated 3 times and assayed on SDS page electrophoresis. Superoxide Dimutase (SOD) activity were assayed by the method of Henary et.al (1976).

Estimation of essential monoterpene oil(s)

Geranium oil estimation was done by steam distillation of 100 g freshly plucked leaves in a clevenger's apparatus (Clevenger, 1928). The oil constituents mainly Geraniol, citronellol and other associated oil contents were determined by Gas liquid chromatography (Perkin –Elemer model 3920 B). The stainless steel column was packed with 10% carbowax (20 mesh) on chromosorb WNAW. Injector and detector temperature was maintained at 200°C. The flow of H₂ was 0.47 cm S⁻¹ data processing for area % was done on a Hewllet Packard integrator model HP-33%.

Statistical Analysis

The results were statistically analyzed for the Least Significant Differences (LSD) using the layout of a complete randomized design (CRD). Further the results were analyzed for the correlation coefficient to determine the relationship among the characters studied, using the relationship $Y = a + b x$.

Results and Discussion

The fresh and dry biomass increased with increase in the level of Zn. (Table 1). Maximum fresh (282.5 g/plant), dry biomass (19.36 g/plant) and leaf area (40.3cm²) was observed at the 0.25 µg Zn ml⁻¹ supply. Growth attributes such as, plant height (64.1 cm) was maximum at 0.50 µg Zn ml⁻¹ supply. Zn levels of 1.0 µg Zn ml⁻¹ were toxic to overall growth parameters. The total chlorophyll content increased upto 0.25 µgZn/ml and then decreased; where as same trend was observed for Chl.a and Chl.b. While Carotenoids were increased from deficient levels of Cr supply to 0.25 µg Zn ml⁻¹ supply. The maximum photosynthetic efficiency in terms of CO₂ exchange rate of 0.82 µgCO₂/m²/s was found at 0.25 µg Zn ml⁻¹ supply.

The saccharide formation was also found to be highest at 0.25 $\mu\text{g Zn ml}^{-1}$ supply. Zn deficiency and Zn toxicity inhibits P_N in cotton (Ohki, 1976), peppermint (Srivastava *et al.*, 1997), Soybean (Ohki, 1978) and sweet mint (Misra et al 2004). A decrease in Chl. photosynthetic pigment content represents a decline in photochemical capacity of leaf at deficient Zn supply. (Ohki, 1976) .

Table 1: Effect of Se acquisition on growth parameters of Geranium.

| Treatment | Plant height (cm) | No. of branches/ plant (Nos.) | Fresh weight /plant (g) | Dry weight/ plant (g) | Leaf area (cm) ² |
|--------------------------------|-------------------|-------------------------------|-------------------------|-----------------------|-----------------------------|
| 0.0 $\mu\text{g Se ml}^{-1}$ | 57.0 | 9 | 218.8 | 14.11 | 8.2 |
| 0.005 $\mu\text{g Se ml}^{-1}$ | 58.0 | 10* | 238.6* | 16.33* | 12.1* |
| 0.05 $\mu\text{g Se ml}^{-1}$ | 61.0* | 13** | 224.8 | 16.81* | 25.2** |
| 0.10 $\mu\text{g Se ml}^{-1}$ | 62.5** | 18** | 252.1** | 17.37** | 39.1** |
| 0.25 $\mu\text{g Se ml}^{-1}$ | 63.4** | 10* | 282.5** | 19.36** | 40.3** |
| 0.50 $\mu\text{g Se ml}^{-1}$ | 64.1** | 10* | 215.5 | 18.46** | 37.2** |
| 1.00 $\mu\text{g Se ml}^{-1}$ | 59 | 8 | 196.2 | 15.85 | 11.2 |
| LSD at 5% | 2.5 | 1.1 | 11.1 | 2.1 | 3.5 |
| 1% | 4.1 | 3.2 | 16.3 | 3.3 | 6.2 |

* ** Values are significantly different at 5% and 1% level.

Peroxidase activity and peroxidase isoenzymes maximum bands were observed at 0.25 $\mu\text{g Zn ml}^{-1}$ cultured plants. The Zn deficient and toxic cultured plants revealed lesser peroxidase activity with lesser peroxidase isoenzymes band profiles. SOD activity showed as Nitro Blue Tetrazolium (NBT) reduction. The maximum SOD (0.248 units/mg.fr.wt.) was obtained at 0.25 $\mu\text{g Zn ml}^{-1}$ cultured plants. In Japanese mint similar report were observed for Mn nutrition (Misra 1996). The maximum monoterpene oil(s) was found at 0.25 $\mu\text{g Zn ml}^{-1}$ treatments. However with respect to oil composition, contents of citronellol, Geraniol, linalool and nerol varied at different Zn treatments (Table 2). As a result of differential Zn supply the content of other micronutrient as Fe, Mn, Zn, and Cu were affected and showed lesser Fe, Mn, Zn and Cu concentrations in shoot tissue of Geranium (Table 3). The maximum (Zn 164 $\mu\text{g g}^{-1}$), Cu (12 $\mu\text{g g}^{-1}$) Mn (98 $\mu\text{g g}^{-1}$ and Fe (537 $\mu\text{g g}^{-1}$) contents were observed at 0.25 $\mu\text{g Zn ml}^{-1}$ treatments.

Table 2: Effect of Se on photosynthetic pigments, photosynthesis (P_N), chlorophyll (Chl) contents and saccharide formation of Geranium.

| Treatment | Chl a mg g ⁻¹ f.wt. | Chl b mg g ⁻¹ f.wt. | Chl a+b mg g ⁻¹ f.wt. | Carotenoids mg g ⁻¹ f.wt. | P _N μg(CO ₂) m ⁻² s ⁻¹ | Saccharides μg (CH ₂ O) m ⁻² s ⁻¹ |
|------------------------------|--------------------------------------|--------------------------------------|----------------------------------------|-----------------------------------------|---------------------------------------------------------------------------|--------------------------------------------------------------------------|
| 0.0 μg Se ml ⁻¹ | 0.68 | 0.50 | 1.18 | 0.086** | 0.15 | 0.102 |
| 0.005 μg Se ml ⁻¹ | 0.79* | 0.56 | 1.35** | 0.051** | 0.19* | 0.129 |
| 0.05 μg Se ml ⁻¹ | 0.94** | 0.61* | 1.55** | 0.024** | 0.75** | 0.510 ** |
| 0.10 μg Se ml ⁻¹ | 1.35** | 0.69** | 2.04** | 0.016** | 0.76** | 0.516** |
| 0.25 μg Se ml ⁻¹ | 1.48** | 0.79** | 2.27** | 0.007** | 0.82** | 0.558** |
| 0.50 μg Se ml ⁻¹ | 1.01** | 0.40 | 1.41** | 0.003* | 0.71** | 0.483** |
| 1.00 μg Se ml ⁻¹ | 0.82* | 0.29 | 1.11 | 0.002 | 0.42** | 0.286* |
| LSD at 5% | 0.11 | 0.08 | 0.07 | 0.0002 | 0.03 | 0.05 |
| 1% | 0.15 | 0.12 | 0.11 | 0.0005 | 0.06 | 0.07 |

* ** Values are significantly different at 1% and 5% level.

Table 3: Effect of Se acquisition on geranium total oil and oil composition.

| Treatments | % of oil | Citronellol % of total oil | Geraniol % of total oil | Linalool % of total oil | Nerol % of total oil |
|------------------------------|----------|-------------------------------|----------------------------|----------------------------|-------------------------|
| 0.0 μg Se ml ⁻¹ | 0.15 | 0.21 | 0.09 | 8.0 | 1.1 |
| 0.005 μg Se ml ⁻¹ | 0.16 | 0.27** | 0.09 | 10.0** | 1.2** |
| 0.05 μg Se ml ⁻¹ | 0.17* | 0.29** | 0.10** | 9.0** | 1.1 |
| 0.10 μg Se ml ⁻¹ | 0.19 | 0.32** | 0.11** | 6.0** | 1.4** |
| 0.25 μg Se ml ⁻¹ | 0.21** | 0.25** | 0.07** | 7.0** | 1.2** |
| 0.50 μg Se ml ⁻¹ | 0.16 | 0.18** | 0.12** | 8.0** | 0.9** |
| 1.00 μg Se ml ⁻¹ | 0.15 | 0.17** | 0.10** | 7.0** | 0.7** |
| LSD at 5% | 0.02 | 0.01 | 0.005 | 0.04 | 0.01 |
| 1% | 0.04 | 0.02 | 0.01 | 0.07 | 0.02 |

* ** Values are significantly different at 5% and 1% level.

Statistical analysis showed a positive significant association between Zn content in leaf and total CO₂ exchange rate (P_N) ($\gamma=0.924 \leq p=0.5\%$) and between P_N and saccharides (sugars) ($\gamma=0.879 \leq p=0.05\%$ content). However, Zn content in leaf was negatively correlated with Chl. a/b ratio. The CER showed a positive significant association with leaf fresh mass ($\gamma = 791 \leq p=0.05\%$), leaf dry mass ($\gamma=692 \leq 0.05\%$), leaf area and total monoterpene oil(s) ($\gamma=0.721 \leq p=0.01$). A positive significant correlation was also observed between saccharides and total oil ($\gamma=0.695 \leq p=0.01\%$). A quadratic trend was observed for all these characters which were comparable in + Zn then the deficient and much higher Zn cultured plants.

Table 4: Effect of Se acquisition on tissue concentrations in shoots of Geranium.

| Treatments | Fe ($\mu\text{g g}^{-1}$) | Mn ($\mu\text{g g}^{-1}$) | Zn ($\mu\text{g g}^{-1}$) | Cu ($\mu\text{g g}^{-1}$) |
|--------------------------------|-----------------------------|-----------------------------|-----------------------------|-----------------------------|
| 0.0 $\mu\text{g Se ml}^{-1}$ | 98 | 26 | 12 | 7 |
| 0.005 $\mu\text{g Se ml}^{-1}$ | 112 | 37** | 19* | 9 |
| 0.05 $\mu\text{g Se ml}^{-1}$ | 142** | 41** | 34** | 11* |
| 0.10 $\mu\text{g Se ml}^{-1}$ | 249** | 57** | 45** | 11 |
| 0.25 $\mu\text{g Se ml}^{-1}$ | 537** | 98** | 64** | 12** |
| 0.50 $\mu\text{g Se ml}^{-1}$ | 419** | 62** | 41** | 7 |
| 1.00 $\mu\text{g Se ml}^{-1}$ | 312** | 53** | 36** | 5 |
| LSD at 5% | 21 | 9 | 7 | 3 |
| 1% | 42 | 11 | 9 | 5 |

* ** Values are significantly different at 5% and 1% level.

The observed maximum changes in CER, photosynthesis pigments content, leaf area, and antioxidant enzymes- peroxidase activities along with other leaf physiological parameters at 0.25 $\mu\text{g Zn ml}^{-1}$ supply indicate that this level is optimum for best growth. At this Zn supply activity of peroxidase and its isoenzymes, the highest CER, chlorophyll content, and saccharides results in maximum flow of photosynthates to the biosynthetic pathways. The main sites of ROS production in the plant cells are the organelles with highly oxidizing metabolic activities or with sustained electron flows: chloroplasts, mitochondria and peroxisomes. (Garczarska et al. 2004). In chloroplasts ROS can be generated by the direct transfer of the excitation energy, or by oxygen reduction in the Mehler reaction (Meloni et al. 2003). In addition, H₂O₂ and O₂⁻ may interact in the presence of certain ions (Co, Al, Cu, Cd, Zn, Se) and Cr chelates to produce highly reactive OH (Sudhakar et al. 2001). In chloroplasts, H₂O₂ is a powerful inhibitor of the Calvin's cycle. Moreover, all ROS can easily react, causing lipid peroxidation and protein denaturation (Yu and Rengel, 1999) therefore the common effect of free radicals are the degradation of the cell membranes (Prochazkova et al. 2001).

To prevent such damages plant cells are equipped with an antioxidative system consisting of the low molecular weight antioxidants such as ascorbate, l-tocopherol, glutathione, Xanthophyll carotenoids and protective enzymes (Donahue et al.1997) and certain essential Zn, Se. and Cr antioxidant elements(Chakmak, 1999).. Thus during the stress metabolic activities, chain reactions resulting from increase in ROS are less terminated the action of SOD, Fe, Mn. Zn and Cu cofactor mediated enzymes, which are major scavenger of singlet oxygen resulting in the formation of H₂O₂ and O₂.(Meloni et al.2003).Less H₂O₂ is then detoxified in the ascorbate – glutathione cycle which involves the oxidation and re-reduction of ascorbate peroxidase and glutathione reductase action, to make water – water cycle(Donahue et al. 1997).Antioxidants is thus oxido-reducible processes(Misra, 2003). In healthy unstressed conditions to the plants not to damage the cell walls and normal physiological activities persists but during the biotic and abiotic stresses of extrinsic factors leads to the antioxidant and antioxidant oxidizing metabolites Further closing of the stomata and stomatal resistance, for utilization of very low concentration of carbon dioxide which enables then to assimilate carbon dioxide even during considerable stomatal closure(El Bassam, 1998). Thus the study was to examine the stresses of environment as well the extrinsic biotic factors to examine simultaneously both ROS of Xanthophyll cycle of carotenoids for dependent energy dissipation and leaf antioxidant system – specially elemental antioxidant chromium(Cr) acquisition in Geranium and its essential monoterpene oil(s) secondary plant products.

The simultaneous reduction in growth, CER content of other micronutrient, and peroxidase activity in leaves indicate the limited availability of photosynthates to essential metabolism and biosynthesis. Utilization of metabolites from primary photosynthetic process into secondary metabolites regulates monoterpene productions (Gershenzon and Croteau, 1991). Thus a close relation between photosynthesis, photorespiration and terpenoid synthesis also exists in essential monoterpene oil(s) bearing plants (Maffei and Codignola 1990). Moreover, the actively growing leaves requires a larger supply of an antioxidants stimulator Zn, in associations with greater supply of photosynthates and since essential oil biosynthesis occurs in these rapidly growing leaves, the initial growth period would require a still greater supply of photosynthates and energy .

Acknowledgement: The authors are thankful to the Director, CIMAP for the encouragement and facilities provided during the studies.

International Journal of Applied Biology and Pharmaceutical Technology Page: 483
Available online at www.ijabpt.com

REFERENCES

- Agrawala, S.C., Sharma, C.P.: The standardization of sand and water culture technique for the study of macro and micronutrients (trace) element deficiencies under Indian conditions. - *Curr. Sci.***40**:424-428, 1961.
- Ajay. Rathore, V.S.: Effect of Zn stress in rice (*Oryza sativa* cv. Manhar) on growth and photosynthetic processes. *Photosynthetica*. - **31**: 571-584, 1995.
- Anonymous,: Wealth of India, Raw Materials, Vol. 7, CSIR, New Delhi, p. 287,1966.
- Anonymous, Annual Report. Horticultural Research Station, Kodaikanal, Tamil Nadu, 1975.
- Anonymous.: Monthly Statistics of the Foreign Trade of India Vol. 11, March,1982.
- Arnon, D.I.:Copper enzymes in isolated chloroplasts. Polyphenol oxidase in *Beta vulgaris*. *Plant Physiol.*- **24**: 1-15,1949.
- Asada,K.: The water-water cycle in chloroplasts : scavenging of active oxygen and dissipation of excess photons .- *Annu Rev Plant Physiol Plant Mol Biol.***50**:601-639,1999.
- Bonet, A.,Poschenrieder, C.,, Barcelo, J.:Chromium III-iron interaction in Fe-deficient and Fe sufficient bean plants.I.Growth and nutrient content.- *J Plant Nutr.* **14**:403-414,1991.
- Chakmak, I.,Engels, C.: Role of mineral nutrients in photosynthesis and yield formation *In: Ringel, Z* ed. *Mineral Nutrition of Crops* New York, USA Haworth Press. Pp. 141-168, 1999.
- Clevenger, J.F.: Apparatus for determination of essential oils. - *J. Amer. Pharmac. Assoc.* - **17**: 346, 1928.
- Dell, B.,Wilson, S.A.: Effect of Zn supply on growth of three species of Eucalyptus seedlings and wheat.- *Plant Soil.* **88**: 377-384,1985.
- Demmig-adams, B., Adams, W W III.: Xanthophyll cycle and light stresss in nature : uniform response to excess direct sunlight among higher plant species.- *Planta.***198**:460-470,1996.
- Douglas, J.S.: Essential oil crops and their uses.- *World Crops.* **21**: 49-54,1969.
- Donahue, J L., Okpodu, C M., Cramer, C L., Grabau, E A., Alscher, R G .: Response of antioxidants to paraquat in Pea leaves.- *Plant Physiol.***113**: 249-257,1997.
- Erickson, R.E.: Industrial importance of monoterpenes and essential oils.- *Lyodia.* **29**: 8-19,1976.
- Gershenzon, J.,Croteau, R.: Regulation of monoterpene biosynthesis in higher Plants. *In: Towers, G.H.N., Safford, H.A.* (ed.): *Biochemistry of the mevalonic Acid pathway to terpenoids*. pp. 99-159. Plenum Press, New York, 1991.
- Havaux, M.,Lutz, C.,grimm B.: Chloroplast membrane photostability in *chlP* transgenic tobacco plants deficient in tocopherols.- *Plant Physiol* **137**:300-310,2003.
- Henry, L., Halliwell, B., Hall, D.:The Superoxide dismutase activity of various photosynthetic organisms measured by a new rapid assay techniques.- *FEBS.***66**: 303.
- Hewitt, E.J.: Sand and water culture methods used in the study of plant nutrition,-Commonwealth Bureau Bot. Plantation Crops Tech. Commun. **22**: 405-439, 1952.
- Hoagland, D.R.,Arnon, D.I.: The water culture method for growing plants without soil.- *Calif. Agric. Exp. Statio .Circ.* **347**: p. 32,1952.
- Huffman Jr, E W., Allaway W H .: Chromium in plants: distribution in tissue, organelles and extracts , and availability of bean leaf Cr to animals. – *J Agric Food Chem.***21**; 982-986, 1973.
- Maffei, M., Codignola, A.: Photosynthesis and herbicide effect on terpene production in peppermint (*Mentha piperita* L.).- *J. Essen. Oil Res.* **2**: 275-286, 1990.
- Marschner, H.: Function of mineral nutrients. Micronutrients *In: Mineral Nutritional of higher plants*. Pp. 269-300, Academic Press, New York, 1986.
- Misra, A.: Genotypic variation of Manganese toxicity and tolerance of Japanese mint.- *J. Herbs, Spices and Medicinal Plants.* **4**: 3-13, 1996.

- Misra, A., Sharma S.: Zn concentration for essential oil yield and menthol concentration of Japanese mint.- Fertilizer Res. **29**: 261-265,1991.
- Misra,A., Srivastava,N.K., Sharma,S.: Role of an antioxidant on net photosynthetic rate,carbon partitioning and oil accumulation in sweet mint.-Agrotropica.15:69-74,2003.
- Munns, R.:Comparative physiology of salt and water stress.-Plant Cell. Environ.**25**:239-250,2002.
- Ohki, K.: Effect of Zn nutrition on photosynthesis and carbonic anhydrase activity in cotton. - Physiol. Plant. **38**: 300-304, 1976.
- Ohki, K.: Zinc concentration of soybean as related to growth, photosynthesis and carbonic anhydrase activity.- Crop. Sci. **18**: 79-82, 1978.
- Poschenrieder, D., Vazquez, M D ., Bonet, A.,Barcelo, J.:Chromium III-iron interaction in iron sufficient and iron deficient bean plants.2. Ultrastructural aspects. J.Plant Nutr.**14**:315-428,1991.
- Pratt, P F .: Chromium.in Diagnostic Criteria for plants and Soils.Ed. H.D. Chapman. Ch. 9, pp. 136-141, University of California,Riverside, 1966.
- Prochazakova, D., Sairam, R K., Srivastava, G C., Singh, D V.:Oxidative stress and antioxidant activity as the basis of senescencein maize leaves.-Plant Sci.**161**:765-771.
- Rajeshwar Rao B.R., Bhattacharya, A.K. : History and botanical nomenclature of rose scented geranium cultivars grown in India.- Indian Perfumer. **36**: 155-161, 1992..
- Ranade, G.S.: Chemistry of Geranium of Geranium oil.-Indian Perfumer. **32**: 61-68, 1988.
- Randall, P.J., Bouma, D.: Zinc deficiency, carbonic anhydrase and photosynthesis in leaves of Spinach.- Plant Physiol. **52**: 229-232,1973.
- Sawa, T., Akaike, T.,Maeda, H.: Tyrosinenitration by peroxinitite formed from nitric oxide and superoxide generated by Xanthine oxidase.- J.Biol.Chem.**275**:32467-32474,2000.
- Sharon, L.M., Kay, E., Lew, J.Y. : Peroxidase isoenzymes from horse raddish roots 1-1 Isolation and Physical properties.- J. Biol. Chem. **241**: 2166-2172,1966.
- Singh, A.: Chemical and biochemical aspects of activated oxygen: singlet oxygen, superoxide anion and related species. In CRC Handbook of free radicals and antioxidants in Biomedicine . Mequel J. Quintanilha, A T., Weber, H. eds. CRC press, inc., Boca Raton, FL, USA. Vol.1 pp. 17-28, 1989.
- Srivastava, N.K., Misra, A., Sharma, S.: Effect of Zn deficiency on net photosynthetic rate, ¹⁴C partitioning, and oil accumulation in leaves of peppermint.- Photosynthetica. **33**: 71-79, 1997.
- Sudhakar, C., Lashmi, A., Giridarakumar, S.:Changes in the antioxidant enzymes efficacy in two high yielding genotypes of mulberry(*Morus alba* L.) under NaCl salinity.- Plant Sci.**161**:613-619,2001.
- Takeda ,T.,Yokota. A.,Shigeoka,S.:Resistance of photosynthesis to hydrogen eroxide in algae,-Plant Cell Physiol.**36**:1089-1095,1995.
- Terry, N.: An analysis of the growth responses in *Beta vulgaris* L. to phototoxic trace elements . II. Chromium. Final report to the Kearney Foundation of Soil Science. July, 1975 - June,1980.
- Verhoeven,A.S.Swanberg,A.,Thao.M.,Whiteman,J.:Seasonal changes in leaf antioxidant systems and xanthophylls cycle characteristics in *Taxus x media* growing in sun and shade environments.-Physiol Plant**123**:428-434,2005.
- Warington, K.: Molybdenum as a factor in the nutrition of lettuce.- Ann Appl Biol **33**: 249-254,1946.
- Yu, Q.,Rengel, Z.: Drought and salinity differentially influence activities of superoxide dismutase in narrow-leafed lupins.- Plant Sci.**142**:1-11.
